

Uniting the Global Gastroenterology Community to Meet the Challenge of Climate Change and Non-Recyclable Waste



Climate change has been described as the biggest global health threat of the 21st century¹ and has significant implications for gastrointestinal (GI) health and disease,² which is the focus of this consensus commentary provided by the World Gastroenterology Organisation (WGO) Climate Change Working Group (CCWG). The CCWG has members from 18 countries representing high, medium, and low-income populations. The WGO includes gastroenterology societies from 108 countries, which represent more than 60,000 medical practitioner members. The CCWG members, who have coauthored this consensus commentary, aim to review the scientific literature on climate and GI health, to encourage education and the undertaking of actionable measures including advocacy, and to further research and collaborations within the global GI community. The CCWG's objective is to assist GI health providers worldwide to adapt to, and mitigate, the effects of climate change on health. The CCWG has partnered with 3 major GI journals, which are copublishing this commentary, given the timeliness and importance of the topic.^{3,4}

Overview of the Impact of Climate Change

Climate Change

The earth's climate is changing because of increasing concentrations of greenhouse gases in the atmosphere, which has already warmed by more than 1°C compared with pre-industrial times. The aim of mitigation efforts is to keep the rise in global mean surface

temperature (GMST) below 2°C by 2100. An increase of 2°C might not seem like a major change, but variations of similar magnitude have been associated with profound alterations of climate in the past. It is important to note that even if this somewhat arbitrary target can be met, catastrophic changes in earth ecosystems can still occur.

The Intergovernmental Panel on Climate Change was established in 1988 with a mandate to assess relevant scientific literature related to climate change. It produces regular reports and updates about climate change. The Intergovernmental Panel on Climate Change projects that as GMST rises, warming will be more pronounced in some areas than others (Figure 1). For example, as global average temperature reaches +2°C, temperature in the Arctic may exceed +5°C. The changes in atmospheric temperature, moisture content, and movement will lead to shifts in rainfall, with decreased or increased precipitation depending on the region (Figure 1). This has implications for access to 2 of the fundamentals for life and health: water and food. Mean temperature change is one metric of temperature change. From a health perspective, other measures, such as the number of days with a maximum temperature greater than the 90th centile are likely more important than the yearly GMST average. Similarly, average changes in precipitation might not be meaningful if rainfall arrives at the wrong time for planting crops. In addition to climate change, other components of the planetary health framework, namely biodiversity loss and toxin exposure, need to be incorporated into considerations of the future of GI and liver health.⁵ Other key parameters of GI and non-GI health include food, water, infrastructure security, biodiversity, changing patterns of GI infections, and migration.²

Biodiversity

Biodiversity is the variety of life, including variations in genes, species, and functional traits. It is more difficult

to link changes in biodiversity, as opposed to direct effects like droughts, to human health because the effects are indirect. The diversity of life on land and ocean is in rapid decline. This is problematic, as greater biodiversity contributes to increasing crop yields, stabilizing fisheries, fodder yield, pest control, plant disease resistance, climate modulation, water purification, and pollination.⁶ Biodiversity loss is associated with increased risk of pathogen transmission,⁷ including viral epidemics.

Pollution, Toxins, and Chemical Exposure

The increase in food production during the last several decades was driven in part by agrochemicals. In fact, two-thirds of global agricultural land is now at risk of pesticide pollution.⁸ These chemicals are found on food for sale, are biologically active in humans, metabolized in part by gut microbiota, and capable of altering gut flora.⁹ There is also a wide variety of biological toxins that are climate sensitive. Mycotoxins are naturally occurring toxins produced by fungi, which have been linked to a variety of illnesses, including hepatocellular carcinoma.¹⁰ Warm, moist, storage environments and consumption of foods of marginal quality increase the risk of exposure. Algal blooms are associated with nitrogen and phosphate contamination of fresh water. Cyanobacteria can produce toxins that are nephrotoxic and hepatotoxic.

Nutrition

In both low- and high-income countries, malnutrition from undernutrition and obesity are simultaneously problematic. More than half a billion people are undernourished. This number will grow as crop yields decline in a warming world, while populations continue to increase.¹¹ Furthermore, rising temperatures and atmospheric CO₂ may decrease the nutritional value of some crops. Decreasing crop yields of lower nutritional value will exacerbate the existing problems of undernutrition,




Figure 1. Maps of mean temperature and precipitation change at +1.5°C and +2.0°C GMST (Global Mean Surface Warming) and the difference between the 2. Greater temperature changes are seen over land masses and in northern latitudes. Bands of decreased precipitation are seen in both southern and northern hemispheres with increased precipitation closer to the poles. The cross hatching highlights areas where at least two thirds of climate models agree on the sign of change as a measure of robustness. From Hoegh-Guldberg et al,¹³ reprinted with permission from the Intergovernmental Panel on Climate Change.

wasting, stunting, and micronutrient deficiency, and paradoxically might drive an increase in obesity, as mild to moderate insecurity has been linked to higher obesity rates,¹² which is likely to contribute to an increase in nonalcoholic fatty liver disease and metabolic-associated fatty liver disease.

Food production contributes as much as 25% of global greenhouse gas emissions. In addition, concerns have been raised regarding the loss of biodiversity from conversion of wilderness to farmland, and the use of pesticides and fertilizers. Attention has also focused on beef and dairy industries because cattle are a source of the potent greenhouse gas methane.

Water

Nearly 40% of the world's 7.9 billion people live in areas with high to

very high water shortages or scarcity. Just as crop yields fall as temperatures rise, so also does access to clean fresh water. Pesticide, chemical use, and contamination with sea water threaten the quality of water available. Utilization of water of questionable quality increases the risk of exposure to toxins and infections.

Rising CO₂ in the atmosphere, the primary driver of climate change, dissolves in the oceans and increases acidity, interfering with the ability of shellfish, coral, and some plankton to form their shells and skeletons. Ocean temperature, chemical change, and overfishing will lead to a decline in this important source of nutrition.

Changing Patterns of Gastrointestinal Disease

The incidence of many GI diseases, such as inflammatory bowel disease

and colorectal cancer, show marked geographic variation, in part due to environmental factors. It can be anticipated that epidemiological shifts in GI disease will occur with a changing environment. There is also a close link between mental health and functional GI disease. Populations are already subject to chronic stress because of struggles to obtain food, water, and shelter. Acute stress from events such as wildfires, floods, and storms, and from forced or elective migration, will add to this mental health burden. An increase in functional GI disorders can be anticipated as climate threats increase.

Two-thirds of the world's megacities are located by the sea. The number of people impacted by sea level rise will depend on the degree of temperature rise. Without adaptation measures, more than 100 million people will be affected by 2050.¹³

Displacement of many of their inhabitants, and the risks from storm surges, will add to the populations forced to migrate because of global warming. Climate change will increase migration as populations move from areas experiencing water and food stress, both from rural to urban areas and between countries. The health needs of these groups and their ability to access care, vary widely depending on their migrant status and the stage of the migrant journey.^{14,15} GI health care providers will need training in diagnosis and management of diseases with which they might not be familiar and in the cultural differences with which different groups approach health care needs.

Diarrheal illnesses consistently rank in the top 5 categories of global disease burden,¹⁶ even before the exacerbating effects of climate change. Transmission of infectious agents to humans depends on pathogen, vector, and host interactions and occurs in a social context. Climate change alters these relationships through changes in temperature, humidity, habitat, access to clean water, food, and sanitation, and is predicted to bring major changes in the epidemiology of infections.¹⁷ Both high and low rainfall extremes are associated with an increased risk of gastrointestinal infections, even in higher-income countries.¹⁸ High rainfall events can lead to contamination of reservoirs with water containing surface organisms and can also overwhelm wastewater treatment. Low rainfall leads to use of water of marginal quality, concentration of pathogens, and reduced sanitation flow. Global warming is also affecting the geographic range of a wide variety of GI infections. One example is the change in the range of *Vibrio cholerae*. Formerly seen in high-income countries rarely, changes in the temperature of the oceans have led to the appearance of cholera along the northwestern seaboard of North America and in the Baltic.¹⁹

The main impact of climate change on the liver will likely be changes in the prevalence of metabolic liver disease secondary to changes in nutrition, increased risk of hepatitis A and E from flooding, changes in the

geographic distribution of other liver infections, such as schistosomiasis, and exposure to toxins, such as aflatoxin.

Acute heat exhaustion has well-described multi-organ effects, including digestive organs. Less is known about the effect of chronically elevated temperatures on intestinal function. A warming atmosphere holds more energy and water. Storms are expected to increase in severity and possibly increase in frequency. GI care is heavily dependent on functioning infrastructure. Storms, wildfires, and floods are already impacting care, and this will likely increase. Planning for adaptive measures to limit the impact of these events should proceed with urgency.

Although the focus of this commentary is on GI health, the health challenges of the changing environment will affect every organ system. Understanding the effects of climate change, biodiversity loss, and toxin exposure on organ systems other than the gut and how that can relate to intestinal function is in evolution.

Carbon Footprint and Nonrecyclable Waste

The health sector and all of us as individuals are significant contributors to greenhouse gas emissions, the majority of which is CO₂, but also includes methane, nitrous oxide, and hydrofluorocarbons. Formation and retention of greenhouse gases in the atmosphere lead to increased energy that translates to global warming due to higher air and water temperatures. The “carbon footprint” represents the CO₂ equivalents that are produced, be it by the burning of fossil and biofuels or use of electricity, heating and cooling, transportation, food, and consumer goods. In the United States, health services account for as much as 8% of total national carbon emissions. The nature of our GI specialty leads to a significant contribution not only to the carbon footprint, but also to huge amounts of nonrecyclable plastics waste.^{20,21}

Plastic waste also comes from many of the nearly 15,000 medical journals, albeit the fraction of online journals is increasing. Many digestive

health and disease journals continue to have plastic covers. The impact of plastic journal covers of 2 journals published by the American Gastroenterological Association alone represented 1.4 metric tons annually (Figure 2A, extrapolated to the journals published by American Gastroenterological Association that are mailed to its members). Of note, journals affiliated with the American Gastroenterological Association and British Society of Gastroenterology have converted the use of plastic covers to labels or to paper covers (Figure 2B and C).

The plastic waste from endoscopic procedures is also difficult to fathom. For example, 1 endoscopic procedure can be estimated to generate 0.54 kg of waste (eg, Figure 2D and E), although the nonrecyclable plastic waste levels become 1.2 kg if suction canisters, tubing, and other accessories are included.²¹ Therefore, for 11 million estimated colonoscopies carried out in the United States,²² the approximate nonrecyclable plastic and nonplastic waste in the United States from colonoscopies alone is nearly 13,200 tons. Similarly, digestive organ endoscopy was reported to be the second highest procedure-associated waste-generating medical department (0.50 kg/procedure) after radiology (0.67 kg/procedure), with plastic surgery being third (0.44 kg/procedure).²³ Notably, plastics can also be degraded to form microplastics, composed of plastic particles <5 mm in size, which can be detected in marine shores and even in human colectomy specimens.²⁴ Specific measures to curtail the pollution impact that is generated by the care provided by gastroenterologists will be discussed.

What Measures Should Be Undertaken by the Gastroenterology Community?

The WGO leadership made the strategic decision to establish a CCWG because of the global impact of climate change as discussed and the need for grassroots efforts. Several medical organizations have begun to form climate change focus groups or consortia, such



Figure 2. Conversion of plastic journal covers to paper, and waste generated by endoscopy procedures. (A) The plastic covers of 2 typical journals weigh 13.6 g; (B, C) Some of the major GI journals have switched from using plastic covers to either paper labels (*Gastroenterology*) or a paper cover (*Gut*). (D) Three bins of endoscopy-generated waste from an endoscopy unit in Melbourne, Australia (for 9 colonoscopies with polypectomies and 1 upper endoscopy, excluding the suction and drainage tubing and canisters). The bins shown in (D) were then weighed after emptying their contents (E). The net weight of waste per procedure was 0.54 kg.

as The Medical Society Consortium, which includes 19 society members.²⁵ We believe that individual GI member societies can act locally and think globally and can be highly effective, given that the needs and resources vary widely across and even within countries. As such, GI societies can be involved directly and indirectly with almost all of the specific efforts, and so can endoscopy centers, GI practices, academic centers, publishers, and individuals (Table 1). In general terms, the efforts revolve around reducing the level of greenhouse gas emissions, curbing nonrecyclable waste, and working toward developing affordable, climate-friendly, substitute disposables.

Measures that can be taken to address the challenges can be adaptive or designed to mitigate. Adaptive measures do not aim to alter the course of climate change but to build resilience. Mitigation efforts are designed to reduce greenhouse gas emissions with an ultimate aim of a carbon neutral environment by 2050.

The initial goals of the WGO CCWG, which we encourage other gastroenterology member societies to establish and undertake, include the following:

1. Advocacy, by publicly supporting efforts of member GI organizations within their countries, be it to support environment-friendly laws or to support increased funding, or to support efforts similar to those that the National Health Service has undertaken.²⁶
2. Education and information, by establishing a website that will house education material and relevant publications; hosting of webinars and podcasts; including climate change sessions at WGO meetings; assembling content and topic interested subgroups of the CCWG and involving other WGO members to develop consensus statements and other in-depth analyses that examine the

health impact of climate change in a continent or region of the world; assess topics such as reusable endoscopy equipment, reducing greenhouse gas emissions from hospitals or institutions, redesigning educational activities to minimize the carbon footprint, including relevant content in the curricula of medical schools; and approach publishers in relation to reducing greenhouse gas emissions in the publication industry.

3. Collaboration with gastroenterology member societies and other medical organizations. One major aspect of this is to work together to decrease the carbon cost of the annual and other sponsored conferences, much of which is related to air travel that could be mitigated by hybrid in-person and remote meetings and other measures.^{27,28} In this regard, the COVID-19 pandemic has already taught us that we

Table 1. What Can Global Gastroenterology Societies, Endoscopy Centers, Practices, Publishers, Individuals and Academic Centers Do to Support Planetary Health?

Endeavor	GI society	Endoscopy center	GI practice	Publisher	Individual	Academic center
Climate change efforts become part of daily life and work activities	✓	✓	✓	✓	✓	✓
Existing endoscopy management committees establish stand-alone (sub)committees to curb waste		✓	✓			
Limit GHG production, primarily the carbon footprint (energy, travel)	✓	✓	✓	✓	✓	✓
Work with endoscopy and supply companies to curb nonrecyclable waste	✓	✓	✓			
Eliminate nonrecyclable plastic waste	✓	✓	✓	✓	✓	✓
Develop and disseminate educational material, and examples of active efforts	✓	✓	✓	✓		✓
Advocate for government and research investments	✓	✓	✓	✓	✓	✓
Provide research support to combat climate change	✓					✓
Embed climate health in undergraduate, graduate and postgraduate curricula						✓

GHG, greenhouse gas.

can very effectively carry out major international virtual meetings. Industry also has an important role to play in helping reduce the carbon footprint of gastroenterologists by reducing waste, exploring the possibilities for reusable equipment, and helping to promote virtual and hybrid conferences.

These steps will contribute to mitigation of the climate crisis through reduction of greenhouse gases and will inform strategies for adaptation to the changes that are underway. For its share, the WGO CCWG will be conducting a survey of its members and developing a strategic plan for WGO to follow within the organization and externally along the lines highlighted above. In addition, the African Middle East Gastroenterology Association is establishing a Climate Action Group, the Canadian Association of Gastroenterology has a special interest section,

and the British Society of Gastroenterology Working Group on Climate Change and Sustainability is undertaking similar strategic planning efforts. It is an opportune moment for the medical community to work together to proactively tackle planetary health challenges as they grow exponentially.

Knowledge Gaps

Research is critically needed in several areas.²⁹ Priorities include the clinical impact of the changing climate and rising temperatures on gastrointestinal health, the relationship between acute and chronic climate stress on functional disorders, the impact on gut microbiota of environmental toxins and decreasing biodiversity, and the role of microplastic exposure on gut health and nutrition. Strategies for endoscopy centers are central to gastroenterology practice,^{29,30} as are development of affordable disposable

plastic substitutes, how best to educate gastrointestinal health care providers, and how to translate knowledge about the climate threat into effective action.

As noted above, climate change has different impacts on health in different regions of the world. The WGO CCWG is setting up subcommittees comprising gastroenterologists from different regions around the world to research and report about the impact of climate change on health in their regions.

Summary

Human progress during the last 100 years came at a huge cost to the environment. As we face the challenge of a warming climate, many measures of GI health are already problematic. The impact of climate change will be felt most severely in lower-income countries, which already struggle to deliver care. Climate change will pose significant additional burdens on digestive health and disease. Meeting

this challenge within our specialty, while providing the best preventive; diagnostic; and therapeutic GI population care, will necessitate a unified global approach to rectify the current impact of climate change and nonrecyclable waste.³¹ We close with quotes from 3 leading world figures, a politician, a cleric, and an 18-year-old activist. “By polluting the oceans, not mitigating CO₂ emissions and destroying our biodiversity, we are killing our planet. Let us face it, there is no planet B” (Emmanuel Macron); “We only have one world. If we destroy it, we’re done for” (Desmond Tutu); and “The eyes of all future generations are upon you. And if you choose to fail us, I say—we will never forgive you” (Greta Thunberg).

DESMOND LEDDIN

Dalhousie University, Canada
Chair, WGO Clinical Research Committee

M. BISHR OMARY

Rutgers University, United States
Past President, American
Gastroenterological Association

ANDREW VEITCH

Royal Wolverhampton Hospitals, United Kingdom
President Elect, British Society of
Gastroenterology

GEOFFREY METZ

Monash University and University of Melbourne, Australia
Chair, Climate Change Working Group,
WGO

NAIMA AMRANI

Mohammed V University, Morocco
President, WGO

LARS AABAKKEN

Oslo University Hospital, Norway

RAJA AFFENDI RAJA ALI

National University of Malaysia, Malaysia

MARIO REIS ALVARES-DA-SILVA

Universidade Federal do Rio Grande do Sul, Brazil

DAVID ARMSTRONG

McMaster University, Canada

SEDAT BOYACIOGLU

Baskent University, Turkey

YE CHEN

Southern Medical University, China

REDA ELWAKIL

Ain Shams University, Egypt

KWONG-MING FOCK

Duke-NUS Medical School and Changi General Hospital, Singapore

SAEED S. HAMID

Aga Khan University, Pakistan

GOVIND MAKHARIA

All India Institute of Medical Sciences, India

FINLAY MACRAE

University of Melbourne, Australia

REZA MALEKZADEH

Tehran University of Medical Sciences, Iran

CHRIS J. MULDER

Amsterdam University Medical Center, the Netherlands

ALEJANDRO PISCOYA

Universidad San Ignacio de Loyola, Peru

MAI LING PERMAN

Fiji National University, Fiji

ANAHITA SADEGHI

Tehran University of Medical Sciences, Iran

ROQUE SÁENZ

Universidad del Desarrollo, Chile

JEAN-CHRISTOPHE SAURIN

Hôpital Edouard Herriot, France

AMNA SUBHAN BUTT

Aga Khan University, Pakistan

KAICHUN WU

Xijing Hospital of Digestive Diseases, China

YEONG YEH LEE

Universiti Sains Malaysia, Malaysia

References

1. Costello A, Abbas M, Allen A, et al. Managing the health effects of climate change: Lancet and University College London Institute for Global Health Commission. *Lancet* 2009;373:1693–1733. Erratum in: *Lancet* 2009;373:2200.
2. Leddin D, Macrae F. Climate change: implications for gastrointestinal health and disease. *J Clin Gastroenterol* 2020;54:393–397.
3. Leddin D, Omary MB, Veitch A, et al. Uniting the global gastroenterology community to meet the

challenge of climate change and non-recyclable waste. *Gut* 2021; 70:2025–2029.

4. Leddin D, Omary MB, Veitch A, et al. Uniting the global gastroenterology community to meet the challenge of climate change and non-recyclable waste. *J Clin Gastroenterol* 2021;55:823–829.
5. Whitmee S, Haines A, Beyrer C, et al. Safeguarding human health in the Anthropocene epoch: report of The Rockefeller Foundation-Lancet Commission on planetary health. *Lancet* 2015;386:1973–2028. Erratum in: *Lancet* 2015;386:1944.
6. Cardinale BJ, Duffy JE, Gonzalez A, et al. Biodiversity loss and its impact on humanity. *Nature* 2012;486:59–67. Erratum in: *Nature* 2012;489:326.
7. Ostfeld RS. Biodiversity loss and the rise of zoonotic pathogens. *Clin Microbiol Infect* 2009;15(Suppl 1):40–43.
8. Tang FHM, Lenzen M, McBratney A, et al. Risk of pesticide pollution at the global scale. *Nat Geosci* 2021;14:206–210.
9. Claus SP, Guillou H, Ellero-Simatos S. The gut microbiota: a major player in the toxicity of environmental pollutants? *NPJ Biofilms Microbiomes* 2016 May 4; 2:16003. Erratum in: *NPJ Biofilms Microbiomes* 2017;3:17001.
10. Bennett JW, Klich M. Mycotoxins. *Clin Microbiol Rev* 2003; 16:497–516.
11. Mbow C, Rosenzweig C, Barioni LG, et al. 2019: Food security. In: Shukla PR, Skea J, Calvo Buendia E, et al., eds. *Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*. In press.
12. Swinburn BA, Kraak VI, Allender S, et al. The global syndemic of obesity, undernutrition, and climate change: the Lancet Commission report. *Lancet* 2019;393:791–846. Erratum in: *Lancet* 2019;393:746.
13. Hoegh-Guldberg O, Jacob D, Taylor M, et al. Impacts of 1.5°C global warming on natural and human systems. In: Masson-Delmotte V, Zhai P, Pörtner H-O, et al, eds.

- Global Warming of 1.5°C. An IPCC Special Report on the Impacts of Global Warming of 1.5°C Above Pre-Industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty. In press.
14. Greenaway C, Castelli F. Migration medicine. *Infect Dis Clin N Am* 2019;33:265–287.
 15. Abubakar I, Aldridge RW, Devakumar D, et al. UCL–Lancet Commission on Migration and Health. The UCL–Lancet Commission on Migration and Health: the health of a world on the move. *Lancet* 2018;392:2606–2654.
 16. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020;396:1204–1222. Erratum in: *Lancet* 2020;396:1562.
 17. Casadevall A. Climate change brings the specter of new infectious diseases. *J Clin Invest* 2020; 130:553–555.
 18. Andrade L, O’Dwyer J, O’Neill E, et al. Surface water flooding, groundwater contamination, and enteric disease in developed countries: a scoping review of connections and consequences. *Environ Pollut* 2018;236:540–549.
 19. Altizer S, Ostfeld RS, Johnson PT, et al. Climate change and infectious diseases: from evidence to a predictive framework. *Science* 2013;341:514.
 20. Williams JA, Kao JY, Omary MB. How can individuals and the GI community reduce climate change? *Gastroenterology* 2020; 158:14–17.
 21. Gayam S. Environmental impact of endoscopy. *Am J Gastroenterol* 2020;115:1931–1932.
 22. Peery AF, Crockett SD, Murphy CC, et al. Burden and cost of gastrointestinal, liver, and pancreatic diseases in the United States: update 2018. *Gastroenterology* 2019;156:254–272.e11.
 23. Vaccari M, Tudor T, Perteghella A. Costs associated with the management of waste from healthcare facilities: an analysis at national and site level. *Waste Manag Res* 2018;36:39–47.
 24. Ibrahim US, Anuar ST, Azmi AA. Detection of microplastics in human colectomy specimens. *JGH Open* 2020;5:116–121.
 25. The Medical Society Consortium on Climate and Health. Member Societies. Available at: medsocietiesforclimatehealth.org. Accessed August 24, 2021.
 26. Delivering a net zero. NHS. Available at: <https://www.england.nhs.uk/greenernhs/wp-content/uploads/sites/51/2020/10/delivering-a-net-zero-national-health-service.pdf>. Accessed August 24, 2021.
 27. Zotova O, Petrin-Cesrosiers C, Gopfert A, et al. Carbon-neutral medical conferences should be the norm. *Lancet Planet Health* 2020; 4:E48–E50.
 28. Leddin D, Galts C, McRobert E, et al. The carbon cost of travel to a medical conference: Modelling the annual meeting of the Canadian Association of Gastroenterology [published online ahead of print July 29, 2021]. *J Can Assoc Gastroenterol* <https://doi.org/10.1093/jcag/gwab021>.
 29. Siddhi S, Dhar A, Sebastian S. Best practices in environmental advocacy and research in endoscopy. *Techn Innov Gastrointest Endosc* in press.
 30. Maurice JB, Siau K, Sebastian S, et al. Green endoscopy: a call for sustainability in the midst of COVID-19. *Green Endoscopy Network. Lancet Gastroenterol Hepatol* 2020;5:636–638.
 31. Smith J, Vignieri S. A devil’s bargain. *Science* 2021;373:34–35.

This article is being published jointly in *Gastroenterology*, *Gut*, and *Journal of Clinical Gastroenterology*. The article is identical except for minor stylistic and spelling differences in keeping with each journal’s style. Citations from any of the 3 journals can be used when citing this article.

Acknowledgments

The authors thank Sarah Williamson and Sunni Losito (American Gastroenterological Association) for assistance with figure preparation, and members of the World Gastroenterology Organisation (WGO) executive committee for reviewing and approving the content of this commentary. The authors also thank Marissa Lopez and Jim Melberg for their support of the WGO Climate Change Working Group, Professor Anthony Capon, Director, Monash University, Sustainable Development Institute, for reviewing our manuscript, and Associate Professor Andrew Metz, Director of Endoscopy, Royal Melbourne Hospital, for kindly providing the images and related information displayed in [Figure 2D](#) and [E](#).

Conflicts of interest

The authors disclose no conflicts.

Most current article

© 2021 by the American Gastroenterological Association, the British Society of Gastroenterology and BMJ Publishing Group, the World Gastroenterology Association, and Wolters Kluwer 0016-5085/\$36.00 <https://doi.org/10.1053/j.gastro.2021.08.001>